



Beef Cattle Handbook



BCH-5460

Product of Extension Beef Cattle Resource Committee
Adapted from the Cattle Producer's Library

Emergency Rations for Wintering Beef Cows

Roger Brownson, Beef Cattle Specialist, Montana State University

Drought conditions greatly reduce the available forage for wintering cattle. During drought periods, both the quality and quantity of hay and winter range forage are often limited.

As a means of stretching limited hay and forage supplies, one pound of grain or other concentrate may be substituted for about two pounds of hay. Six pounds of grain is the practical limit for replacing hay in the daily ration for wintering a 1,100 pound cow. Feeding six pounds of grain will save 12 pounds of hay per head per day. This is a substantial saving over a full feed of hay at prevailing prices.

Corn silage may replace part or all of the hay in the ration at the rate of three pounds of silage to one pound of hay.

Tables 1 and 2 list examples of possible rations substituting grain for part of the forage and using more straw in the ration. Lesser amounts of grain can be used if hay is more plentiful or relative prices change.

If cows are thin at calving time, conception rates will be reduced 60 - 90 days later at breeding time. After calving, cows' daily nutrient requirements increase 30 - 50 percent because of the lactation requirement.

A high-phosphorus mineral mixture and salt should be available to cows at all times.

Proposed rations are based on making maximum use of grain in meeting the minimum standards for wintering beef cows. A deficiency in vitamin A in some rations may be met by feeding one pound of a commercial protein supplement fortified with at least 10,000 I.U. of vitamin A or one pound of dehydrated alfalfa pellets. Rations are based on cattle being allowed access to

range forage or confined to a drylot. The ration assumes that cattle on poor range will have a daily intake of ten pounds of forage and cattle on very poor range a daily intake of about five pounds.

These proposed rations are general in nature, and adjustments may be needed to meet an individual rancher's needs. In drought conditions, all cows should receive 20,000 - 30,000 I.U. of vitamin A during the last 90 days of pregnancy to ensure against vitamin A deficiencies and subsequent losses. Feed manufacturers can provide the needed level of vitamin A in a variety of protein supplements or in a custom pellet. Green color in feed is an indication of carotene or vitamin A.

Cattle can store vitamin A in their liver and fat from 4 - 6 months. Cattle on dry pasture during the major part of the summer and on cured hay and straw during the winter can become deficient by spring. That, coupled with the cow's greater need in late pregnancy and lactation, could cause serious problems if the need is not met.

Table 1. Rations for 1,000 Pound Pregnant Cow.

Feed	Daily feed (lb.)	Protein (lb.)	Energy (ME Mcal)	Ca (g)	P (g)	Vitamin A (1,000s I.U.)
Dairy requirement	19.6	1.6	17.3	23	18	25
#1 Barley, oats, or wheat straw	11.0	0.3	7.4	7	3	0
Alfalfa hay	11.0	1.6	10.4	60	10	280
Total	22.0	1.9	17.8	67	13	280
#2 Barley, oats, or wheat straw	12.0	0.4	8.0	8	4	0
Corn	5.0	0.4	7.5	1	5	1.8
Protein supplement (P 20%, Ca 1%, Ph 1%)	2.0	0.4	2.8	9	9	?
Total	19.0	1.21	18.3	18	18	1.8 ²
#3 Barley, oats, or wheat straw	12.0	0.4	7.4	7	3	0
Corn silage	30.0	0.7	11.4	12	9	25
Protein supplement (P 20%, ph 1%)	1.0	0.2	1.4	9	4	?
Total	43.0	1.3	20.2	19	16	25
#4 Barley, oats, or wheat straw	12.0	0.4	7.4	7	3	0
Barley	6.0	0.7	8.1	2	11	0
Protein supplement (P 20%, Ph 1%)	1.0	0.2	1.4	9	4	?
Total	19.0	1.31	16.91	18	18	0 ¹

¹ Marginal unless additional consumption occurs.² Add mineral supplement and/or vitamin A as needed.**Table 2. Rations for 1,100 Pound Cows Nursing Calves (Average Milking Ability).**

Feed (lb)	Daily feed (lb)	Protein (ME Mcal)	Energy (g)	Ca (g)	P (1,000s I.U.)	Vitamin A
Dairy requirement	21.6	2.0	19.9	27	22	38
#1 Barley, oats, or wheat straw	10.0	0.3	6.7	6	3	0
Alfalfa hay	14.0	2.1	13.3	76	12	357
Total	24.0	2.4	20.0	82	151	357
#2 Barley, oats, or wheat straw	12.0	0.4	7.4	7	3	0
Barley	7.0	0.8	16.3	2	13	0
Protein supplement (P 20%, Ca 1%, Ph 1%)	2.0	0.8	2.8	9	9	?
Total	21.0	2.0	26.5	182	25	02
#3 Barley, oats, or wheat straw	12.0	0.4	7.4	7	3	0
Corn silage	40.0	0.9	15.6	16	12	33
Protein supplement P 20%, ph 1%)	2.0	0.8	2.8	9	9	?
Total	54.0	2.1	25.8	32	24	33

¹ Add a phosphorus supplement or phosphorus in the salt.² Add a mineral supplement and/or vitamin A as needed.

Authors:

Roger Brownson, Beef Cattle Specialist, Montana State University

Adapted from the Cattle Producer's Library

This publication was prepared in cooperation with the Extension Beef Cattle Resource Committee and its member states and produced in an electronic format by the University of Wisconsin-Extension, Cooperative Extension. Issued in furtherance of Cooperative Extension work, ACTS of May 8 and June 30, 1914.

BCH-5460 Emergency Rations for Wintering Beef Cows